AUTHOR INDEX OF VOLUME 36

Abdul Sattar, M. H. and Bahamish, H. S. Occurrence of Fusarium wilt on potato in Democratic Yemen, 403
Abisgold, J. D. and Fishpool, L. D. C. A method for estimating the population sizes of whitefly nymphs (*Bemisia tabaci* Genn.) on cassava, 287
Adu, J. K. *see* Salako, E. A., 380
Ahmad, S. *see* Shafi, M. M., 371
Ahmed, S. M. *see* Shafi, M. M., 371
Alhubaishi, A. A. *see* Walkey, D. G. A., 195
Altieri, M. A. *see* Gold, C. S., 362
Aneja, K. R. and Srinivas, B. Leaf-spot disease of Field bindweed, *Convolvulus arvensis* – a new disease record, 75
Aneja, K. R. and Srinvas, B. Leaf spot disease of water hyacinth, *Eichhornia crassipes* – a new disease record from India, 405
Aneja, K. R., Srinivas, B. and Singh, K. Three new pathogenic fungi of waterhyacinth from India, 76
Armstrong, J. W. *see* Hansen, J. D., 359
Arny, D. C. *see* Sah, D. N., 154
Ashraf, M. *see* Mahmood, T., 282
Awoderu, V. A. Yield loss attributable to neck-rot of rice caused by *Pyricularia oryzae* Cav. in Cote d'Ivoire, 394

Bahamish, H. S. *see* Abdul Sattar, M. H., 403
Banaras, M. *see* Mahmood, T., 282
Bashir, N. H. H. *see* Yassin, K. M. 230
Bellotti, A. C. *see* Gold, C. S., 362
Bhardwaj, S. P. and Panwar, K. S. Outbreak of *Heliothis armigera* (Hubner) and *Plusia orichalcea* (F.) on Black Zira in the dry temperate region of India, 73
Bibo, J. *see* Hawton, D., 147
Biliwa, A. *see* von Berg, A., 270
Brayford, D. *see* Waller, J. M., 181
Buctuanon, E. M. *see* Morallo-Rejesus, B., 114

Calumpang, S. M. F. *see* Tejada, A. W., 237
Chandrasena, J. P. N. R. Torpedograss (*Panicum repens* L.) control with lower rates of glyphosphate, 336
Charaya, M. U. *see* Tyagi, S., 384
Chindo, P. S. and Kahn, F. A. Control of root-knot nematodes, *Meloidogyne* spp. on tomato, *Lycopersicon esculentum* Mill., with poultry manure, 332
Cooper, J. F. *see* Harris, E. G., 162
Cordero-Jenkins, L. E., Jiron, L. F. and Lezama, H. J. Notes on the biology and ecology of *Ptecticus testaceus* (Diptera: Stratiomyidae) a soilder fly associated with mango fruit (*M. indica*) in Costa Rica, 285
Cowie, R. H. *see* Tiben, A., 296

Dabek, A. J. and Waller, J. M. Black leaf streak and viral leaf streak: new banana diseases in East Africa, 157
DeFrank, J. The response of nine protea species to spray applications of fluazifop-p, 145
Dube, V. P. *see* Tyagi, S., 384
Dumaliang, N. *see* Schreiner, I. H., 93
Dutt, A. K. and Saha, J. L. Insect pest management of winter planted cotton in coastal rice-fallows of West Bengal, 89
Duveiller, E., Notteghem, J. L., Rott, P., Snacken, F. and Maraite, H. Bacterial sheath brown rot of rice caused by *Pseudomonas fuscovaginae* in Malagasy, 151

Easton, C. S. *see* Gunning, R. V., 293
Egwuatu, R. I. *see* Emehute, J. K. U., 279
Elliot, C. C. H. *see* Parker, J. D., 234
Emehute, J. K. U. and Egwuatu, R. I. Effects of field populations of cassava mealybug, *Phenacoccus manihoti*, on cassava yield and *Epidinocarsis lopezi* at different planting dates in Nigeria, 279
Evangelista, C. C. and Tangonan, N. G. Reaction of 31 non-senescent sorghum genotypes to stalk rot complex in southern Philippines, 214

Fishpool, L. D. C. *see* Abisgold, J. D., 287
Flower, L. S. *see* Harris, E. G., 162
Fomba, S. N. and Singh, N. Crop losses caused by brown spot disease in mangrove swamps of northwestern Sierra Leone, 387
Forrester, N. W. *see* Gunning, R. V., 293

Gabriel, T. Pest Management, Women and rural extension, 173
Gadalla, B. H. *see* Yassin, K. M., 230
Ghewande, M. P. Biological control of groundnut (*Arachis hypogaea* L.) rust (*Puccinia arachidis* Speg.) in India, 17
Giga, D. P. and Mazarura, U. M. Malathion resistance in *Sitophilus zeamais* (Mostch) in Zimbabwe, 320
Gold, C. S., Altieri, M. A. and Bellotti, A. C. Effects of intercropping and varietal mixtures on the cassava hornworm, *Erinnyis ello* L. (Lepidoptera: Sphingidae), and the stemborer, *Chilomima clarkei* (Amsel) (Lepidoptera: Pyralidae), in Colombia, 362
Govindarajulu, V. *see* Peter, C., 312
Greenland, D. J. Better crop protection information, 220
Greenup, L. R. *see* Gunning, R. V., 293
Gunasekera, S. A. *see* Shanthichandra, W. K. N., 375
Gunasekera, S. A., Shanthichandra, W. K. N. and Price, T. V. Disease incidence, severity and pod yield of winged bean *Psophocarpus tetragonolobus* accessions and *Psophocarpus scandens*, 207
Gunning, R. V., Forrester, N. W., Easton, C. S. and Greenup, L. R. Relationship between DDT and pyrethroid resistance in *Heliothis armigera* (Hubner) (Lepidoptera: Noctuidae) in Australia, 293

Gupta, Alamelu, *see* Parihar, N. S., 15
Gupta, M. P. *see* Ram, Shri., 258

Haciwa, H. C. *see* Kannaiyan, J., 21
Hamm, R. T. *see* Lim, T. K., 23
Hansen, J. D. and Armstrong, J. W. The failure of field sanitation to reduce infestation by the mango weevil, *Cryptorhynchus mangiferae* (F.) (Coleoptera: Curculionidae), 359
Harris, E. G., Cooper, J. F., Flower, L. S., Smith, S. C. and Turner, C. R. Toxicity of insecticide aerosol drops to tsetse flies. I. Some effects of temperature, formulation and drop size, 162
Harvey, G. L. *see* Hawton, D., 147
Haseeb, A. and Pandey, R. Root-knot nematodes-a constraint to cultivation of Davana, *Artemisia pallens*, 317
Hawton, D., Johnson, I. D. G., Loch, D. S., Harvey, G. L., Marley, J. M. T., Hazard, W. H. L., Bibo, J. and Walker, S. R. A guide to the susceptibility of some tropical crop and pasture weeds and the tolerance of some crop legumes to several herbicides, 147
Hazard, W. H. L. *see* Hawton, D., 147
Heath, R. R. *see* Jansson, R. K., 263
Hedge, R. K. *see* Narayana Bhat, M., 73
Hennessey, R. D., Neuenschwander, P. and Muaka, T. Spread and current distribution of the cassava mealybug, *Phenacoccus manihote* (*Homoptera: Pseudococcidae*), in Zaire, 103
Hiremath, I. G., Vastard, S. and Patil, S. J. *Lamprosema charesalis* (Walker) (Lepidoptera: Pyraustidae), a new pest of tumeric, *Curcuma domestica* Val. in India, 74

Imeokparia, P. O. Performance of a new selective rice herbicide in a lowland rice field in Nigeria, 343

Jansson, R. K., Proshold, F. I., Mason, L. J., Heath, R. R. and Lecrone, S. H. Monitoring sweetpotatoe weevil (Coleoptera: Curculionidae) with sex pheromone: effects of dosage and age of septa, 263
Javaid, I. Application of insecticides on cotton in Zambia: timing of spray applications, 1
Jeremiah, H. *see* Parker, J. D., 234
Jindal, K. K. and Thind, B. S. Control of bacterial blight of cowpea (*Vigna unquiculata* (L.) Walp.), 211
Jiron, L. F. *see* Cordero-Jenkins, L. E., 285
Johnson, I. D. G. *see* Hawton, D., 147

Kabissa, J. C. B. Seasonal occurrence and damage by *Pectinphora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) to cotton in Eastern Tanzania, 356
Kahn, A., Host status of commonly grown cultivars of chillies (*Capsicum frutescens*) and sweet pepper (*C. annuum*) to the root-knot nematode (*Meloidogyne incognita*) in northern Nigeria, 329
Kahn, F. A. *see* Chindo, P. S., 332
Kambal, M. A. *see* Tiben, A., 296

Kannaiyan, J. and Haciwa, H. C. Economic benefits of spraying fungicides to control groundnut foliar diseases in Zambia, 21
Kanwar, B. B. *see* Raj, Desh. 10
Keshava, Bhat, S., Vidyasagar, P. S. P. V. and Sujatha, A. Record of *Rattus rattus wroughtoni* as a pest of oil palm seedlings, 75
Key, G. Pre-harvest crop losses to the African striped ground squirrel, *Xerus erythropus*, in Kenya, 223
Khokhar, K. M. *see* Mahmood, T., 282

La Croix, E. A. S. Studies on macadamia nut borer (*Crytophlebia* spp.) in northern Malawi, 97
Latis, T. A note on the carpenter moth *Salagena* sp., a new pest of cashew in Zambia, 276
Lecrone, S. H. *see* Jansson, R. K., 263
Lezama, H. J. *see* Cordero-Jenkins, L. E., 285
Lim, T. K., Hamm, R. T. and Mohamad, R. B. Persistency and volatile behaviour of selected chemicals in treated soil against three basidiomycetous root disease pathogens, 23
Loch, D. S. *see* Hawton, D., 147

MacFarlane, J. H. Damage assessment and yield losses in sorghum due to the stem borer *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae) in Northern Nigeria., 131
Magallona, E. D. *see* Tejeda, A. W., 237
Magambo, M. J. S. *see* Onsando, J. M., 346
Mahmood, T., Khokhar, K. M. M. Banaras., and Ashraf, M. Effect of environmental factors on the density of leaf hopper, *Amrasca devastan* (distant) on okra, 282
Maraite, H. *see* Duveiller, E., 151
Maredia, K. M. and Mihm, J. A. Damage by southwestern corn borer (*Diatraea grandiosella* Dyar) on resistant and susceptible maize at three plant growth stages in Mexico, 141
Marley, J. M. T. *see* Hawton, D., 147
Mason, L. J. *see* Jansson, R. K., 263
Matthews, G. A. Changes in application techniques used by the small scale cotton farmer in Africa, 166
Matthews, G. A. *see* Omar, D., 159
Mazarura, U. M. *see* Giga, D. P., 320
Mbata, G. N. Suitability of maize varieties for the oviposition and development of *Plodia interpunctella* (Hubner) (Lepidoptera: Pyralidae), 122
Mihm, J. A. *see* Maredia, K. M., 141
Mohamad, R. B. *see* Lim, T. K., 23
Morallo-Rejesus, B., Buctuanon, E. M., and Rejesus, R. S., Defining the economic threshold determinants for the Asian corn borer, *Ostrinia furnacalis* (Guenee) in the Philippines, 114
Mote, U. N. and Murthy, D. K. Studies of loss estimation and relative susceptibility of genotypes of sorghum to earhead caterpillar (*Heliothis armigera* Huebner), 108
Muaka, T. *see* Hennessey, R. D., 103
Murthy, D. K. *see* Mote, U. N., 108

Nafus, D. M. *see* Schreiner, I. H., 93
Narayana Bhat, M. and Hegde, R. K. Occurrence of *Curvularia verruculosa* in *Lagerstroemia lanceolata* in India, 73
Neuenschwander, P. *see* Hennessey, R. D., 103
Notteghem, J. L. *see* Duveiller, E., 151
Nsemwa, L. T. H. and Shao, F. M. New record of plant pathogenic fungi in Tanzania, 74
Nwanze, Kanayo, F. and Sivakumar, M. V. K. Insect pests of pearl millet in Sahelian West Africa-11. *Raghuva albipunctella* De Joannis (Noctuidae, Lepidoptera): distribution, population dynamics and assessment of crop damage, 59
Nyambo, Brigitte, T. Effect of natural enemies on the cotton bollworm, *Heliothis armigera* Hübner (Lepidoptera: Noctuidae) in Western Tanzania, 50

Ogeda, K. *see* Oloo, G. W., 224
Ogunwolu, E. O. Damage to cowpea by the legume pod borer, *Maruca testulalis* Geyer, as influenced by infestation density in Nigeria, 138
Oladiran, A. O. The effects of fungicides and insecticides singly and in combination on the control of brown blotch, pod borer infestation and yields of *Vigna unguiculata* (L) Walp, 397
Oloo, G. W. and Ogeda, K. The incidence of *Chilo partellus* (Swinh.) (Pyralidae) and the contribution of natural enemies to its mortality under intercropping system in Kenya, 244
Omar, D. and Matthews, G. A. The rainfastness of permethrin deposits of emulsifiable concentrate (EC) and ultra-low-volume (ULV) formulation on brussels sprout leaves, 159
Omolo, J. G. *see* Onsando, J. M., 346
Onsando, J. M., Magambo, M. J. S., and Omolo, J. G. Evaluation of Basta on Kikuyu grass (*Pennisetum clandestinum* chiov.), in Kenya, 346

Palmer, J. M. Identification of the common thrips of Tropical Africa (Thysanoptera: Insecta), 27
Pandey, R. *see* Haseeb, A., 317
Panwar, K. S. *see* Bhardwaj, S. P., 73
Parihar, N. S. and Gupta, Alamelu. Dissipation of endosulfan in green foliage, pods and grains of red gram (*Cajanus cajan* L.), 15
Parker, J. D., Uk, S., Elliot, C. C. H. and Jeremiah, H. Potential health hazards from the consumption of quelea sprayed with fenthion avicide, 234
Patil, S. J. *see* Hiremith, I. G., 74
Pearce, M. J. A new trap for collecting termites and assessing their foraging activity, 310
Pearce, M. J. *see* Tiben, A., 296
Pervez, A. *see* Shafi, M. M., 371
Peter, C. and Govindarajulu, V. Management of blossom thrips, *Megalurothrips usitatus* on pigeonpea, 312
Price, T. V. *see* Gunasekera, S. A., 207
Price, T. V. *see* Shanthichandra, W. K. N., 375
Proshold, F. I. *see* Jansson, R. K., 263

Raj, Desh. and Kanwar, B. B. Minimizing insecticide use against cauliflower pests in India, 10
Ram, Shri and Gupta, M. P. Integrated pest management in lucerne (*Medicago sativa* L.) and its economics in India, 258
Rejesus, R. S. *see* Morallo-Rejesus, B., 114
Rott, P. *see* Duveiller, E., 151

Sah, D. N. and Arny, D. C. Susceptibility of maize germplasm to bacterial stalk rot caused by *Erwinia chrysanthemi* pv. zeae, 154
Sah, R. P. *see* Sharma, S. B., 327
Saha, J. L. *see* Dutt, A. K., 89
Salako, E. A. and Adu, J. K. Effect of foliar fungicides on foliar diseases and pod, haulm and plant crude protein yields of groundnut, 380
Salman, A. G. and Sayed, A. A. Construction techniques for the control of the sand termite, *Psammotermes hybostoma* Desneux in traditional housing in Egypt, 68
Sarr, E. *see* Sharma, S. B., 71
Sayed, A. A. *see* Salman, A. G., 68
Schreiner, I. H., Nafus, D. M. and Dumaliang, N. Growth and survival of the Asian corn borer *Ostrinia furnacalis* Guenee (Lep: Pyralidae) on alternative hosts in Guam, 93
Serit, M., and Tan Keng Hong. Immature life table of a natural population of *Dacus dorsalis* in a village ecosystem, 305
Shafi, M. M., Pervez, A., Ahmad, S. and Ahmed, S. M. Role of some taste additives to enhance poison bait acceptance in the black rat, *Rattus rattus* L., 371
Shanthichandra, W. K. N. *see* Gunasekera, S. A., 207
Shanthichandra, W. K. N., Gunasekera, S. A. and Price, T. V. Diseases and pests of the winged bean (*Psophocarpus tetragonolobus*) [L] DC) in Sri Lanka, 375
Shao, F. M. *see* Nsemwa, L. T. H., 74
Sharma, S. B., Sah, R. P., Singh, Onkar, and van Rheenen, H. A. Root-knot nematode disease of chickpea in Nepal, 327
Sharma, S. B., Subrahmanyam, P. and Sarr, E. Plant parasitic nematodes associated with groundnut in Niger, 71
Shukla, R. P. Spatial distribution of key pests of mustard and their natural enemies in India, 85
Singal, S. K. and Singh, Z. Studies of plant oils as surface protectants against pulse beetle, *Callosobruchus chinensis*

(L.) in chickpea, *Cicer arietinum* L. in India, 314
Singal, S. K. and Toky, O. P. Carry over of bruchid, *Caryedon serratus* (Olivier) (Coleoptera) from field to stores through seeds of *Acacia nilotica* (L.) Willd, in India, 66
Singh, K. *see* Aneja, K. R., 76
Singh, N. *see* Fomba, S. N., 387
Singh, Onkar, *see* Sharma, S. B., 327
Singh, Z. *see* Singal, S. K., 314
Sivakumar, M. V. K. *see* Nwanze, Kanayo, F., 59
Smith, S. C. *see* Harris, E. G., 162
Snacken, F. *see* Duveiller, E., 151
Srinivas, B. *see* Aneja, K. R., 75
Srinivas, B. *see* Aneja, K. R., 76
Srinivas, B. *see* Aneja, K. R., 405
Stary, P. *see* Völkl, W., 249
Stechmann, D. H. *see* Völkl, W., 249
Subrahmayam, P. *see* Sharma, S.B., 71
Sudoi, V. Evaluation of different acaricides for control of red crevice mite *Brevipalpus phoenicis* Geijskes (Acari-Tenuipalpidae) infesting tea, 349
Sujatha, A. *see* Keshava Bhat, S., 75

Tan Keng Hong *see* Serit, M.
Tangonan, N. G. *see* Evangelista, C. C., 214
Tejada, A. W., Calumpang, S. M. F. and Magallona, E. D. The fate of carbofuran in rice-fish and livestock farming, 237
Thind, B. S. *see* Jindal, K. K., 211
Tiben, A., Pearce, M. J., Wood, T. G., Kambal, M. A. and Cowie, R. H. Damage to crops by *Microtermes najdensis* (Isoptera, Macrotermitinae) in irrigated semi-desert areas of the Red Sea Coast. 2. Cotton in the Tokar Delta region of Sudan, 296
Toky, O. P. *see* Singal, S. K., 66
Torreno, H. S. Parasitization behaviour and efficiency of the Braconid, *Microgaster manilae* (Ashmead), against the cutworm, *Spodoptera litura* (F.), 128
Turner, C. R. *see* Harris, E. G., 162
Tyagi, S., Dube, V. P. and Charaya, M. U. Biological control of the purple blotch of onion caused by *Alternaria porri* (Ellis) Ciferri, 384

Uk, S. *see* Parker, J. D., 234

van Rheenen, H. *see* Sharma, S. B., 327
Varma, B. K. Plant quarantine inspection, procedures and facilities for the import and export of seeds and vegetative propagules at the International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), 216
Vastard, S. *see* Hiremith, I. G., 74
Vidyasagar, P. S. P. V. *see* Keshava Bhat, S., 75
Völkl, W., Stechmann, D. H. and Stary, P. Suitability of five species of Aphidiidae (Hymenoptera) for the biological control of the banana aphid *Pentalonia nigronervosa* Coq. (Homoptera, Aphididae) in the South Pacific, 249
von Berg, A. and Biliwa, A. Control of the larger corn borer (*Prostephanus truncatus*) (Horn) (Coleoptera; Bostrichidae) in traditional maize storage structures in southern Togo, 270

Waite, G. K. *Amblypelta* spp. (Hemiptera: Coreidae) and green fruit drop in lychees, 353
Walker, S. R. *see* Hawton, D., 147
Walkey, D. G. A., Alhubaishi, A. A. and Webb, M. J. W. Plant virus diseases in the Yemen Arab Republic, 195
Waller, J. M. and Brayford, D. Fusarium diseases in the tropics, 181
Waller, J. M. *see* Dabek, A. J., 157
Webb, M. J. W. *see* Walkey, D. G. A., 195
Wood, T. G. *see* Tiben, A., 296

Xiao-Shui, W. *Altica cyanae* (Col: Chrysomelidae) for the biological control of *Ludwigia prostrata* (Onagraceae) in China, 368

Yassin, K. M., Bashir, N. H. H. and Gadalla, B. H. Effects of endosulphan, chlorpyrifos and their mixtures on *Bemisia tabaci* of Sudan Gezira, 230